

# Reconciling estimates of the contemporary North American carbon balance among terrestrial biosphere models, atmospheric inversions, and a new approach for estimating net ecosystem exchange from inventory-based data<sup>1</sup>

DANIEL J. HAYES\*<sup>1</sup>, DAVID P. TURNER<sup>†</sup>, GRAHAM STINSON<sup>‡</sup>, A. DAVID MCGUIRE<sup>§</sup>, YAXING WEI\*, TRISTRAM O. WEST<sup>¶</sup>, LINDA S. HEATH<sup>|||</sup>, BERNARDUS DEJONG<sup>\*\*</sup>, BRIAN G. MCCONKEY<sup>††</sup>, RICHARD A. BIRDSEY<sup>‡‡</sup>, WERNER A. KURZ<sup>‡</sup>, ANDREW R. JACOBSON<sup>§§</sup>, DEBORAH N. HUNTZINGER<sup>¶¶</sup>, YUDE PAN<sup>‡‡</sup>, W. MAC POST\* and ROBERT B. COOK\*

\*Environmental Sciences Division, Oak Ridge National Laboratory, Oak Ridge, TN 37831, USA, <sup>†</sup>Department of Forest Ecosystems and Society, Oregon State University, Corvallis, OR 97331, USA, <sup>‡</sup>Pacific Forestry Centre, Canadian Forest Service, Victoria, BC V8Z 1M5, Canada, <sup>§</sup>U.S. Geological Survey, Alaska Cooperative Fish and Wildlife Research Unit, University of Alaska Fairbanks, Fairbanks, AK 99775, USA, <sup>¶</sup>Joint Global Change Research Institute, Pacific Northwest National Laboratory, College Park, MD 20740, USA, <sup>|||</sup>USDA Forest Service, Durham, NH 03824, USA, <sup>\*\*</sup>El Colegio de la Frontera Sur (ECOSUR), Villahermosa, C.P. 86280, Tabasco, Mexico, <sup>††</sup>Agriculture and Agri-Food Canada, Ottawa, ON K1A 0C5, Canada, <sup>‡‡</sup>USDA Forest Service, Newtown Square, PA 19073, USA, <sup>§§</sup>NOAA Earth System Research Lab, Boulder, CO 80305, USA, <sup>¶¶</sup>School of Earth Sciences & Environmental Sustainability, Northern Arizona University, Flagstaff, AZ 86011, USA, <sup>|||</sup>Currently on secondment to the Global Environment Facility, Washington, DC 20006, USA

## Abstract

We develop an approach for estimating net ecosystem exchange (NEE) using inventory-based information over North America (NA) for a recent 7-year period (ca. 2000–2006). The approach notably retains information on the spatial distribution of NEE, or the vertical exchange between land and atmosphere of all non-fossil fuel sources and sinks of CO<sub>2</sub>, while accounting for lateral transfers of forest and crop products as well as their eventual emissions. The total NEE estimate of a  $-327 \pm 252$  TgC yr<sup>-1</sup> sink for NA was driven primarily by CO<sub>2</sub> uptake in the Forest Lands sector ( $-248$  TgC yr<sup>-1</sup>), largely in the Northwest and Southeast regions of the US, and in the Crop Lands sector ( $-297$  TgC yr<sup>-1</sup>), predominantly in the Midwest US states. These sinks are counteracted by the carbon source estimated for the Other Lands sector ( $+218$  TgC yr<sup>-1</sup>), where much of the forest and crop products are assumed to be returned to the atmosphere (through livestock and human consumption). The ecosystems of Mexico are estimated to be a small net source ( $+18$  TgC yr<sup>-1</sup>) due to land use change between 1993 and 2002. We compare these inventory-based estimates with results from a suite of terrestrial biosphere and atmospheric inversion models, where the mean continental-scale NEE estimate for each ensemble is  $-511$  TgC yr<sup>-1</sup> and  $-931$  TgC yr<sup>-1</sup>, respectively. In the modeling approaches, all sectors, including Other Lands, were generally estimated to be a carbon sink, driven in part by assumed CO<sub>2</sub> fertilization and/or lack of consideration of carbon sources from disturbances and product emissions. Additional fluxes not measured by the inventories, although highly uncertain, could add an additional  $-239$  TgC yr<sup>-1</sup> to the inventory-based NA sink estimate, thus suggesting some convergence with the modeling approaches.

**Keywords:** agriculture, carbon cycle, climate change, CO<sub>2</sub> emissions, CO<sub>2</sub> sinks, forests, inventory, modeling, North America

Received 7 November 2011; revised version received 7 November 2011 and accepted 21 November 2011

Correspondence: D.J. Hayes, tel. + 865 574 7322, fax + 865 241 3685, e-mail: hayesdj@ornl.gov

<sup>1</sup>This manuscript has been authored by UT-Battelle, LLC, under Contract No. DE-AC05-00OR22725 with the U.S. Department of Energy. The publisher, by accepting the article for publication, acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this manuscript, or allow others to do so, for United States Government purposes.

## Introduction

North American ecosystems have had a significant influence on the global carbon budget by acting as a large sink of atmospheric CO<sub>2</sub> in recent decades (Fan *et al.*, 1998; Myneni *et al.*, 2001; Butler *et al.*, 2010). Although the exact contribution is uncertain, analyses of the global C budget suggest that this North

American terrestrial sink may be responsible for nearly a third of the combined global land and ocean sink of atmospheric CO<sub>2</sub> (Pacala *et al.*, 2007). A recent review of late 20th Century carbon balance estimates for terrestrial ecosystems in North America (NA) compiled for the State of the Carbon Cycle Report (SOCCR) found a wide range of results, with estimates of the magnitude of the continental-scale CO<sub>2</sub> sink extending between 0.1 and 2.0 PgC yr<sup>-1</sup> (King *et al.*, 2007), although the terrestrial sink based on inventories reported in this document was 0.5 PgC yr<sup>-1</sup> with uncertainty of about 50%<sup>1</sup> (Pacala *et al.*, 2007). By comparison, fossil fuel emissions over NA (from Canada, the US and Mexico combined) in the early 21st Century are estimated to be approximately 1.8 PgC yr<sup>-1</sup> (Boden *et al.*, 2010).

Although fossil fuel emissions are calculated with relatively high precision, understanding the fate of those emissions with respect to sequestration in terrestrial ecosystems requires data and methods that can reduce uncertainties in the diagnosis of land-based CO<sub>2</sub> sinks. The wide range in the land surface flux estimates is related to a number of factors, but most generally because of the different methodologies used to develop estimates of carbon stocks and flux, and the uncertainties inherent in each approach. The alternative approaches to estimating continental scale carbon fluxes that we explored herein can be broadly classified as applying a *top-down* or *bottom-up* perspective. *Top-down* approaches calculate land-atmosphere carbon fluxes based on atmospheric budgets and inverse modeling. *Bottom-up* approaches rely primarily on measurements of carbon stock changes (the 'inventory' approach) or on spatially distributed simulations of carbon stocks and/or fluxes using process-based modeling (the 'forward model' approach).

Atmospheric inversion models (AIMs) infer surface fluxes by reference to a sample of atmospheric CO<sub>2</sub> concentration (mixing ratio) measurements coupled with models of surface flux and atmospheric transport (Gurney *et al.*, 2002; Ciais *et al.*, 2010). These inverse analyses provide constraints on estimates of land-atmosphere carbon exchange at a detailed temporal resolution, relying on the strong diurnal and seasonal cycles in CO<sub>2</sub> concentration in the observations. However, these estimates are associated with large uncertainties from the limited density of observation networks, uncertainty in the transport models, and errors in the inversion process (Gurney *et al.*, 2004; Baker *et al.*, 2006). Further, AIMs typically operate at a coarse spatial resolution and provide limited detail on the processes controlling the carbon sources and sinks.

Biomass inventories provide valuable constraints on changes in the size of carbon pools over years to decades (e.g. Pacala *et al.*, 2001; Peylin *et al.*, 2005). Invento-

ries are designed to precisely measure standing stocks in forests on longer time scales, and to estimate and analyze the dynamics of growth, harvest, and mortality. However, the inventory measurement approach can only detect measurable changes in vegetation which usually occurs over a number of years, and therefore re-measurements in most inventory programs are taken periodically. There is a high likelihood that dynamics and fluxes will be under-sampled or missed altogether; for instance, inventory sampling can produce reliable estimates of biomass, but other carbon pools (e.g. litter and soil C stocks) are not sampled at the same intensity in all areas. Inventory-based modeling can be used to estimate growth and disturbance impacts, but does not yet provide full capability in partitioning the forcing brought about by non-disturbance factors (Stinson *et al.*, 2011). On the other hand, inventory and commerce data sets can often be used to quantify the storage, emissions and/or lateral movement of carbon in product pools, which are typically not well-characterized in modeling approaches.

The forward model approach builds from understanding the underlying processes controlling carbon dynamics and can be used to simulate the dynamics of multiple ecosystem components through a class of models referred to as terrestrial biosphere models (TBMs). However, TBMs contain substantial uncertainty due to the sheer number of often poorly understood underlying processes simulated. They also vary widely in the data used to drive them, in the particular processes simulated, and in their level of detail (Schwalm *et al.*, 2010, Huntzinger *et al.*, in press). Yet, TBMs simulate the impacts of multiple driving forces and controlling mechanisms of land-atmosphere CO<sub>2</sub> exchange, incorporate non-linear system behaviors, make predictions at spatial and temporal scales relevant to global and regional carbon cycles, and allow for exploration of the impacts of underlying processes.

Each of the three general approaches (inventory, forward and inverse modeling) build on different knowledge foundations and employ different driver data. A suite of results on NA ecosystem carbon flux from extant model simulations (based on both TBMs and AIMs) have been organized by the North American Carbon Program (NACP; Denning, 2005; Wofsy and Harris, 2002) under the regional and continental interim-synthesis (RCIS) activities (Huntzinger *et al.*, in press). The RCIS activities focus on 'off-the-shelf' model simulations and other recently published studies as a pre-cursor to more formal model inter-comparison activities. Here, we assembled and analyzed available inventory-based data on NA ecosystem carbon cycle components as an additional perspective alongside the forward and inverse approaches avail-

able from the RCIS. We developed novel techniques for comparison of the inventory-based data against results from the TBMs and AIMS at common spatio-temporal scales and flux indicators.

## Materials and methods

The magnitude of carbon sources and sinks is defined as the vertical exchange of CO<sub>2</sub> between the surface (land or ocean) and the atmosphere, hereafter referred to as net ecosystem exchange (NEE). In this analysis, we used estimates of NEE for the biosphere where fossil fuel emissions are excluded from the calculation. From the land perspective, NEE is primarily the balance between CO<sub>2</sub> uptake in vegetation through net primary production (NPP) and release via the heterotrophic respiration (R<sub>h</sub>) of dead organic matter, plus emissions from fires and the decay of harvested forest and agricultural products (Chapin *et al.*, 2006). Here we used the sign convention from an atmospheric reference point whereby a negative value of NEE represents land surface uptake (a sink) and a positive value represents CO<sub>2</sub> emissions to the atmosphere (a source).

The geographic domain of this study included the three countries of NA (Canada, the US, and Mexico) and the reference time period was approximately 2000–2006. NEE estimates were made at an annual time step and considered lateral in addition to vertical transfers of carbon. Spatial scale became important where a relatively large amount of carbon is transported laterally (as harvested biomass products transferred off-site or as dissolved carbon transported in rivers, for example). In these cases, the CO<sub>2</sub> was considered a sink at the location where it was taken up, but became a source at the location where it was eventually returned to the atmosphere (through product decay or in-stream decomposition, for example). In this analysis, carbon flux was estimated at the scale allowable by the various inventory-based data sets (i.e., by inventory ‘reporting zones’). We distinguished three sectors (Forest Lands, Crop Lands, and ‘Other’ Lands) within 97 spatial units (total number of ‘reporting zones’ across the three countries) in each (Table 1). The 97 ‘reporting zones’ refer to the sum of US states, Canadian managed ecoregions, and Mexican states for which inventory data were available. The carbon flux estimates from 7 inverse and 17 forward models were compiled from those submitted to the NACP-RCIS activity (<http://nacarbon.org/nacp>; Huntzinger *et al.*, in press). Here we focused on ecosystem carbon fluxes, whereas fossil fuel emissions are discussed for comparison but were not included in the budgets.

### *Inventory-based estimates of NEE*

For the national-level reporting of greenhouse gas (GHG) inventories in the context of the Framework Convention on Climate Change (FCCC), the protocol is generally to track changes in pool sizes using data collected or modeled for carbon pools of different key land-based sectors, such as forest and agricultural lands along with other non-forest (e.g., grasslands), settled (developed and built-up) lands, and areas of land use change (Parson *et al.*, 1992). In this study, we compiled GHG inventory-based data on productivity, ecosystem

carbon stock change and harvested product stock change for managed Forest Lands and Crop Lands in Canada and the United States. Additional information was used to fill in data on carbon balance in Other Lands, including data on human and livestock use/consumption of harvested products. For Mexico, our analysis accounted primarily for carbon flux due to land use change. Data on carbon exchange for each sector were summarized by reporting zone, with spatial and temporal coverage of the data sets noted in Table 1a and details on methods by country and sector described in the Supporting Information.

The conceptual model used to organize the various sector-specific data sets is illustrated by Fig. 1. The data for both the Forest Lands and Crop Lands sectors (left side of diagram) were based upon estimated stock changes within the vegetation and soil carbon pools. According to the conceptual model, all the stock changes in these pools represented vertical exchange of CO<sub>2</sub> with the atmosphere (i.e., NEE) except for (1) the vertical exchange of non-CO<sub>2</sub> trace gases, (2) the leaching of carbon from the system via river export and (3) the ‘lateral’ movement of carbon between sectors and reporting zones. Lateral movement occurs via changes in land use as well as the harvest and transport of forest and agricultural products. Where available, data on these fluxes were used to produce more precise estimates of NEE for each sector in each reporting zone from the stock change information. Total average annual NEE (NEE<sub>TOT</sub>) is the combination of NEE estimated for the Forest Lands (NEE<sub>F</sub>), Crop Lands (NEE<sub>C</sub>) and Other Lands (NEE<sub>O</sub>) sectors for each reporting zone:

$$NEE_{TOT} = NEE_F + NEE_C + NEE_O. \quad (1)$$

Which and how the underlying component fluxes, and their inventory-based data sources, were used to estimate NEE<sub>F</sub>, NEE<sub>C</sub>, and NEE<sub>O</sub> are described in the sections below. Note that, in the equations given, not only NEE but also all component flux values were treated with the atmospheric reference sign convention whereby a negative value represents a CO<sub>2</sub> sink effect and a positive value a source effect of that component. By this definition, fire emissions have positive values, harvest removals have negative, and positive values of stock change represent losses in different C pools and vice versa.

### *Forest lands sector inventories*

Although the equations differ depending on the data source, our calculations of NEE<sub>F</sub> were, in general, based on inventory estimates of stock changes adjusted for the lateral transfer of harvest removals:

$$NEE_F = \Delta Live + \Delta DOM + H_R + H_E. \quad (2)$$

The change in C stocks in live biomass ( $\Delta Live$ ) included overstory trees, understory vegetation and roots, whereas change in dead organic matter stocks ( $\Delta DOM$ ) included dead trees, down woody debris, litter and soil organic carbon pools. Carbon removed in wood harvest ( $H_R$ ) was considered as a sink from the stand where to wood was grown. However, an additional variable was calculated to represent the proportion

**Table 1** (a) Characteristics (temporal coverage, spatial coverage, variables included, literature reference) of inventory-based estimates used in this study

Data type/ Name	Temporal coverage	Spatial coverage	Variables included in NEE	Reference
Canada managed forest	2000–2006	( <i>n</i> = 15)	NPP, Rh, Fire(CO <sub>2</sub> ), Harvest	Kurz <i>et al.</i> (2009), Stinson <i>et al.</i> (2011)
Canada agriculture	2000–2006 (avg)	Harvest area ( <i>n</i> = 15)	Harvest, ΔDOM	Environment Canada (2011)
Canada 'Other'	2000–2006	( <i>n</i> = 15)	Stock changes	EPA (2011)
	2006	( <i>n</i> = 15)	Livestock emissions	Environment Canada (2011)
US forest	2000–2006 (avg)	Forest area ( <i>n</i> = 49)*	ΔLive, ΔDOM, harvest	Heath <i>et al.</i> (2011), Smith <i>et al.</i> (2009)
US cropland	2000–2006	Cropland Area ( <i>n</i> = 48)	Harvest, ΔDOM	West <i>et al.</i> (2011)
US other	2000–2006	Grasslands, Settlements ( <i>n</i> = 50)*†	Stock changes	EPA (2011)
	2006	( <i>n</i> = 50)	Livestock emissions	EPA (2011)
	2000–2006	( <i>n</i> = 50)	Human respiration	West <i>et al.</i> (2009)
Mexico	1993–2002 (avg)	Mexico ( <i>n</i> = 32)	Stock changes (LUC), forest harvest, forest biomass increment	deJong <i>et al.</i> (2010)

(b) Characteristics (temporal coverage, spatial coverage, variables included, literature reference) of inverse model estimates used in this study

Data type/Name	Temporal coverage	Spatial coverage	Reference
CarbonTracker	2000–2007	North America ( <i>n</i> = 97)	Peters <i>et al.</i> (2007)
Jena	2001–2007	North America ( <i>n</i> = 97)	Rodenbeck <i>et al.</i> (2003)
LSCE-1	2000–2004	North America ( <i>n</i> = 97)	Peylin <i>et al.</i> (2005)
LSCE-2	2000–2006	North America ( <i>n</i> = 97)	Chevallier <i>et al.</i> (2007)
MLEF-PCTM	2003–2004	North America ( <i>n</i> = 97)	Butler <i>et al.</i> (2010)
U. Michigan	2000–2001	North America ( <i>n</i> = 97)	Michalak <i>et al.</i> (2004)
U. Toronto	2000–2003	North America ( <i>n</i> = 97)	Deng <i>et al.</i> (2007)

(c) Characteristics (temporal coverage, spatial coverage, variables included, literature reference) of terrestrial biosphere model estimates used in this study

Data type/Name	Temporal coverage	Spatial coverage	Variables included in NEE	Land use (LU) & disturbance	Reference
Diagnostic (MODIS)					
MOD17+	2000–2004	North America ( <i>n</i> = 97)	Re-GPP		Reichstein <i>et al.</i> (2005)
EC-MOD	2000–2006	North America ( <i>n</i> = 97)	NEE <sup>‡</sup>		Xiao <i>et al.</i> (2008)
Diagnostic (AVHRR)					
SiB3	2000–2005	North America ( <i>n</i> = 97)	Re-GPP		Baker <i>et al.</i> (2008)
CASA	2002–2003	North America ( <i>n</i> = 97)	Re-GPP		Randerson <i>et al.</i> (1997)
CASA GFEDv2	2000–2005	North America ( <i>n</i> = 97)	Re-GPP + Fire	Prescribed fire	van der Werf <i>et al.</i> (2006)
CLM-CASA'	2000–2004	North America ( <i>n</i> = 97)	(Ra + Rh)–GPP		Randerson <i>et al.</i> (2009)
Prognostic					
CLM-CN	2000–2004	North America ( <i>n</i> = 97)	(Ra + Rh)–GPP + Fire	Prescribed LU, prognostic fire	Thornton <i>et al.</i> (2009)
DLEM	2000–2005	North America ( <i>n</i> = 97)	(Ra + Rh)–GPP + Fire + Prod	Prescribed LU, harvest, fire, storms	Tian <i>et al.</i> (2011)
CanIBIS	2000–2005	US & Canada ( <i>n</i> = 66)	(Ra + Rh)–GPP	Prescribed fire	Kucharik <i>et al.</i> (2000)
ISAM	2000–2005	North America ( <i>n</i> = 97)	(Ra + Rh)–GPP	Prescribed LU	Yang <i>et al.</i> (2009)
LPJmL	2000–2005	North America ( <i>n</i> = 97)	(Ra + Rh)–GPP + Fire	Prescribed fire	Bondeau <i>et al.</i> (2007)

Table 1 (continued)

(c) Characteristics (temporal coverage, spatial coverage, variables included, literature reference) of terrestrial biosphere model estimates used in this study

Data type/Name	Temporal coverage	Spatial coverage	Variables included in NEE	Land use (LU) & disturbance	Reference
MC1	2000–2006	Continental US ( $n = 49$ )	(Ra + Rh)–GPP + Fire	Prescribed LU, prognostic harvest & fire	Bachelet <i>et al.</i> (2003)
BEPS	2000–2004	North America ( $n = 97$ )	(Ra + Rh)–GPP		Ju <i>et al.</i> (2006)
ORCHIDEE	2001–2005	North America ( $n = 97$ )	(Ra + Rh)–GPP + Fire	Prescribed LU, prognostic harvest & fire	Krinner <i>et al.</i> (2005)
TEM6	2000–2006	North of 45°N ( $n = 14$ )	(Ra + Rh)–GPP + Fire + Prod	Prescribed LU, harvest, fire	Hayes <i>et al.</i> (2011)
VEGAS2	2000–2005	North America ( $n = 97$ )	(Ra + Rh)–GPP + Fire	Prognostic fire	Zeng <i>et al.</i> (2005)

\*includes Alaska.

†includes the District of Columbia.

‡NEE (and GPP) are empirically derived from MODIS variables.




Fig. 1 Conceptual diagram of the continental-scale carbon budget, including the land-atmosphere exchange of CO<sub>2</sub> (NEE), based on data available from the inventory-based approaches that estimate carbon stock changes, fluxes and transfers among forest, crop, and other lands.

of  $H_R$  that was emitted during the processing of harvested wood into products ( $H_E$ ). This processing, or ‘primary consumption’, was assumed to occur largely at the mill, and so we allocated this source term within the Forest Lands sector of the reporting zone in which the wood was harvested. The remainder (i.e.,  $H_R - H_E$ ) was assumed to be transported off-site and added to the national-level forest product pool that resides in the Other Lands sector (described below).

The data set on forest carbon accounting in Canada’s Managed Forest Area used here employed the ‘stock-plus-flow’ approach, which starts with data from a compiled set of inventories and then models the components of change. Flux data were produced using the Carbon Budget Model of the

Canadian Forest Sector (CBM-CFS3), which uses stand-level growth data to estimate annual carbon uptake along with detailed annual natural disturbance (e.g., fire, insects) and harvest data to track carbon transfers through the system (Kurz *et al.*, 2009; Stinson *et al.*, 2011). Natural disturbance and harvest removals data were from various provincial-level reporting sources in Canada (Stinson *et al.*, 2011). The stock change terms ( $\Delta\text{Live} + \Delta\text{DOM}$ ) as shown in Eqn (2) also included non-CO<sub>2</sub>/non-vertical exchanges and these fluxes were separated out of the NEE<sub>F</sub> calculation. These more detailed component fluxes were estimated by CBM-CFS3, and so NEE<sub>F</sub> for Canada was calculated from the available indicator variables as:

$$NEE_F = \Delta\text{Live} + \Delta\text{DOM} - (\text{FireC} - \text{FireCO}_2) + H_R + H_E \quad (3)$$

where the carbon remaining in harvested products after primary consumption (i.e.,  $H_R - H_E$ ) and the non- $\text{CO}_2$  component of fire emissions (i.e.,  $\text{FireC} - \text{FireCO}_2$ ) were excluded from the vertical flux component of the overall stock change. For Canada, we used 30% as the proportion of  $H_R$  emitted in primary consumption, based on an analysis of 2010 FAO statistics (FAOStat; <http://faostat.fao.org/>) and Canadian harvest data for the period 2000–2006. Therefore:  $H_E$  is equal to  $0.3 \times H_R$  for each reporting zone.

The forest inventory data sets for the US were based on the forest surveys of the U.S. Department of Agriculture (USDA) Forest Service's Forest Inventory and Analysis (FIA) program (Bechtold & Patterson, 2005). These estimates were coupled with carbon expansion factors (Bechtold & Patterson, 2005; Smith *et al.*, 2006; Heath *et al.*, 2011) and estimates of carbon stock changes were derived from the Carbon Calculation Tool (CCT; Smith *et al.*, 2010), which is used to produce the GHG inventory for US forest lands in the UNFCCC reports (EPA, 2011). Harvest removals ( $H_R$ ) were from published US Forest Service data sets (Smith *et al.*, 2009). Estimates of the proportion of  $H_R$  emitted in primary consumption ( $H_E$ ) were provided by Smith *et al.* (2006), who showed that the proportion lost within the first year following harvest (which we assumed occurs primarily at the mill) ranges from 20% to 40% across species group and region in the US. As such, we used 30% as a representative emissions (from primary consumption) fraction, which is the same as that used for the Canada data set. State-level data on fire emissions from US forests were not available for the time period of this study; however, in terms of our NEE calculation, fire emissions were implicit in the total stock change (i.e. fire emissions would have accumulated as biomass had there been no fire) and considered a source of carbon to the atmosphere. The US forest data represents *net* stock change, meaning that fluxes stemming from land use change (LUC; i.e. forest land area converted to other land use, and other land converted to forest land) were also implicit (i.e. integrated in) in the stock change data. The corresponding change in carbon stocks directly attributed to fire and LUC cannot be explicitly separated from the total stock change. Therefore,  $NEE_F$  for the US Forest Lands sector used exactly that as shown in Eqn (2), without the modification for non- $\text{CO}_2$  fire emissions as used in Canada.

As with the Canada forest data set, the Mexico inventory data can be described as being based on the 'stock-plus-flow' approach. For Mexican forests, the data set was based on a carbon accounting methodology in which mean carbon stock density by forest type was distributed according the areal extent of each type at an initial point in time, and stock change was estimated according to the biomass increment (growth) and harvest amount in managed forests, and area of forest conversion over a subsequent period of time. Using this methodology, the study by deJong *et al.* (2010) calculated for the 1993–2002 time period: (1) biomass losses resulting from the conversion of forests to other land use ( $\Delta\text{Live}_{\text{LUC}}$ ); (2) the associated change in soil carbon stocks resulting from LUC ( $\Delta\text{Soil}_{\text{LUC}}$ ); (3) carbon uptake due to the regrowth of forests on

abandoned agricultural or other lands ( $\Delta\text{Live}_{\text{ABND}}$ ); and (4) the net carbon balance between uptake (growth) and emissions (harvest) in managed forests ( $\Delta\text{Live}_{\text{MNGD}}$ ). Fire emissions were included with respect to burning in forest conversion, but the reporting methodology does not take into account fire emissions or other natural carbon fluxes (growth, mortality) from unmanaged land.  $NEE_F$  was calculated by summing the four average annual stock change components from the study by deJong *et al.* (2010):

$$NEE_F = \Delta\text{Live}_{\text{LUC}} + \Delta\text{Soil}_{\text{LUC}} + \Delta\text{Live}_{\text{ABND}} + \Delta\text{Live}_{\text{MNGD}}. \quad (4)$$

For this study, we distributed the magnitude of each component flux proportionately by an estimate of the relative area of each LU/LC class contained in each state, as described in the Supporting Information. Without more detailed data, we assumed that commercial harvest and fuelwood harvest occurred proportional to the relative area of each forest type.

### Crop lands sector inventories

To estimate NEE for croplands for this study, we collected estimates of crop productivity (NPP), harvest ( $H_R$ ) and changes in soil carbon stocks ( $\Delta\text{Soil}$ ) over the 2000–2006 time period for Canada (Environment Canada, 2011) and the US (West *et al.*, 2011). The detail regarding the source and methodologies used in the crop inventories are provided in the Supporting Information as well as by those references cited.  $NEE_C$  was calculated for each reporting zone in Canada and the US as:

$$NEE_C = \Delta\text{Soil} + H_R, \quad (5)$$

where all crop harvest removals (i.e.,  $H_R$ ) were considered a Crop Lands sector sink in the reporting zone where they were harvested; unlike the treatment of harvested wood products, we assumed no primary consumption emissions within the Crop Lands sector. We considered  $\Delta\text{Live}$  in croplands to be equal to zero on an annual basis since the assumption of the data was that NPP is equal to the crop harvest plus residue. We then assumed that, within the same year, the residue carbon was returned to the atmosphere (via combustion or decomposition) or incorporated into the soil C pool.

Data specific to crop productivity and harvest in Mexico were not available for this study, and croplands were not mapped separate from other agricultural lands and forest plantations in the study by deJong *et al.* (2010). As such, we were not able to report estimates of sources and sinks for the Mexican cropland sector separately in this study, but rather included the contribution of soil carbon stock changes from agricultural establishment and abandonment in the Other Lands sector for Mexico.

### Other lands sector inventories

The Other Lands sector was used in this study to include two additional fluxes: (1) net surface carbon fluxes from lands not included in Forest Land or Crop Land sectors (i.e. grasslands, settlements and other lands) and (2)  $\text{CO}_2$  emissions from the

combustion, decay, and respiration of carbon in harvested forest and crop products.  $NEE_O$  was calculated for Canada and the US by combining various component fluxes according to the following equation:

$$NEE_O = NEE_G + NEE_S + E_H + E_L + E_F, \quad (6)$$

which considered the net carbon balance of grassland areas ( $NEE_G$ ), the net carbon balance of human settlement areas ( $NEE_S$ ),  $CO_2$  emissions from human respiration ( $E_H$ ),  $CO_2$  emissions from livestock respiration ( $E_L$ ) and  $CO_2$  emissions from the decay of harvested forest products ( $E_F$ ). For  $NEE_G$  and  $NEE_S$  we used general, area-weighted estimates of 'Grassland' and per-capita estimates of 'Settlements/Other' sink categories reported in the EPA GHG inventory for years 2000–2006 (EPA, 2011). We then extrapolated area-weighted  $NEE_G$  and per-capita  $NEE_S$  according to the area or human population represented by each category in each reporting zone. The area of Other Lands in each reporting zone of Canada and the US is calculated as the remainder of the total area of each zone after subtracting the Forest Land and Crop Land areas from the inventory data sets. The estimates of the product emission terms ( $E_H + E_L + E_F$ ) are described in the next section and in the Supporting Information.

The data set containing state-level estimates of carbon flux from the Other Lands sector in Mexico was developed using the same Eqn (4) as the Forest Lands sector. To calculate  $NEE_O$  for Mexico, we included the component flux estimates for the non-forest types of the LU/LC classification used by deJong *et al.* (2010), which included agricultural lands, forest plantations, scrubland, grassland, wetland, and other non-forest classes. Fuelwood harvest was calculated as a sink in the Forest Lands sector, with emissions transferred to the Other Lands sector (in the same reporting zone that the fuelwood was harvested). The area represented by the Other Lands sector in each reporting zone of Mexico was calculated as the remainder of the total area of each zone after subtracting the forest class areas based on the LU/LC categories used by deJong *et al.* (2010).

#### *Lateral transfer and emissions of harvested products*

In this analysis, the key to linking the Forest Lands and Crop Lands sectors with the Other Lands sector was through data on harvested products (both forest and agricultural), thereby allowing for tracking the movement of carbon between sector and reporting zone. Here, we used the 'atmospheric flow' approach that, according to IPCC Guidelines, accounts for net emissions or removals within national – or, in our case, reporting zone – boundaries (Eggleston *et al.*, 2006). Carbon removal due to growth and emissions due to primary consumption were accounted for in the Forest Land or Crop Land sector of the 'producing' zone. The carbon emissions from secondary consumption were attributed to the Other Lands sector, redistributed proportionately among the reporting zones of the relevant country according to simple assumptions about where the products are likely to be consumed (and thus where the carbon there-in will be returned to the atmosphere as  $CO_2$ ). Our accounting reflects the assumption that some amount of

the carbon in harvested products was not likely to be emitted directly from within the sector (Forest Lands or Crop Lands) that it originates from, but rather in the 'other' lands that the consumers (i.e., humans and livestock) occupy.

Harvested product emissions occurred via the combustion, decay and respiration of harvested wood products (HWP) and harvested crop products (HCP) through secondary consumption by humans (HWP and HCP) and livestock (HCP). Based on the forest and crop inventory data sets, harvested products were summed to national-level pools and adjusted for international imports and exports. Foreign trade of HWP was determined from the FAOStat database for Canada and the Environmental Protection Agency (EPA) GHG Inventory (EPA, 2011) for the US. Foreign trade of HCP was based on the Canadian Socio-Economic Information Management System (Statistics Canada) and the USDA Economic Research Service's 'Foreign Agricultural Trade of the United States' 2010 report. Our simple assumption for allocating the trade-adjusted remaining pools was based on distributing product emissions to the level of the reporting zones proportionally according to human population (HWP and HCP) and data on livestock emissions (HCP). The national-level total HCP from this study was allocated to both human and livestock consumption. The human portion was calculated based on per-capita consumption and emissions (West *et al.*, 2009). The remaining HCP was then allocated to livestock emissions (i.e. assuming no net annual storage of HCP) considering emissions factors for different species, rather than population counts directly.  $CO_2$  emissions from livestock consumption of HCP were distributed proportional to year 2006 methane emissions through enteric fermentation per reporting zone for the US from the USDA Greenhouse Gas Inventory (2008) and for Canada from the Statistics Canada (2006) Census of Agriculture. In the case of longer lived HWP, we used data on stock change in national wood product pools (EPA, 2011) to account for both carbon storage and emissions. Since wood products can be longer lived than our study period, the product pools included 'inherited' stocks and emissions from wood products harvested prior to our study period. Details for the collection and analysis of HWP and HCP carbon data and flux estimates are provided in the Supporting Information.

#### *Uncertainty in inventories and additional fluxes*

We characterized the uncertainty of the inventory-based estimates of NEE presented herein by attaching previous analyses of the major components of the carbon budget of each sector considered in this study (Table S11). We represented the uncertainty around each component in relative terms (as% of the estimate) based on the relevant Monte-Carlo analysis reported in national-level GHG inventories, where available, as well as expert judgment based on previous studies. The ranges of uncertainty on the sector-level mean estimates were calculated by summing the upper and lower bounds for each component flux of the sector; the percent uncertainty, then, was the range between the bounds relative to the mean total flux estimate of the sector.

With respect to the aggregate estimate of continental-scale NEE, another major source of uncertainty came from those components of the carbon budget that are potentially important, but were not measured or estimated by the GHG inventories. These components included fluxes from unmanaged/not inventoried lands (wetlands), potentially important mechanisms not captured (woody encroachment on non-forest landscapes), other potential carbon storage pools (rivers and reservoirs) and lateral fluxes (dissolved organic carbon export from soil through rivers to the ocean) not measured in the inventories (Table S12). The 'best estimate' flux for each of these components was reported in the SOCCR (Chapter 3; Pacala *et al.*, 2007), where expert judgment suggests that these estimates are essentially 100% uncertain.


### *Inventory and model data comparison*

To compare flux estimates at the national and sub-national scales, we included here results based on the inverse modeling approach from among the suite of NACP-participating AIMS that submitted surface flux estimates at  $1 \times 1$  degree grid cell resolution to the RCIS activity. The models within this set of seven (Table 1b) differ in their various formulations and methodologies, including the spatial/temporal resolution, the land model for generating the *a priori* surface fluxes, and the atmospheric transport model employed in the inversion. In two cases (Peters *et al.*, 2007; Lokupitiya *et al.*, 2008), emissions from biomass burning were prescribed and the reported NEE is the sum of the residual land flux (done by inversion) and the prescribed biomass burning flux.

We included in this study a set of 17 NACP-participating TBMs that contributed regional or continental scale results of

recent-era (~2000 to 2006) simulations based on the forward modeling approach. All models were required to submit their best estimate of NEE, which included different component fluxes depending on the particular model (Table 1c). Most models contributed results that cover all the reporting zones for NA used in this study ( $n = 97$ ), whereas some models (CanIBIS, MC1, TEM6) covered subsets of the region. The individual models were based on different simplifying assumptions, used different environmental driving data and initial conditions, and formulated the processes controlling carbon exchange in different ways. Most broadly they were differentiated into prognostic models, which are self-regulating with respect to leaf area index, and diagnostic models in which leaf area (or a surrogate) is prescribed from remote sensing imagery. Among the prognostic models there were significant differences with respect to treatment of fire and other disturbances. Details of these model differences are described by Schwalm *et al.* (2010) and Huntzinger *et al.* (in press).

The contributed results from TBMs and AIMS for the NACP-RCIS were standardized to monthly flux estimates at  $1 \times 1$  degree resolution over the NA land area. To allow comparison at the temporal and spatial scales of the inventories, monthly data were first aggregated to annual flux estimates. These annual flux estimates were then translated from the  $1 \times 1$  degree grid to an estimate for each sector within each reporting zone (Fig. 2). The map of reporting zones consisted of 97 analysis polygons that matched the resolution of the GHG inventory-based data, as described above. The coverage of sectors (Forest Lands, Crop Lands, and Other Lands) was based on a  $1 \text{ km}^2$  grid using aggregation of land cover classes from the GLC2000 data set (Bartholome & Belward, 2005). Juxtaposing these data layers permitted the TBMs and AIMS



**Fig. 2** Forest and cropland reclassification for model-data processing of country/sector carbon flux estimates. The left panel shows the spatial distribution of forest, crops and 'other' lands as per our categorization of the GLC2000 map product. The right panel shows the results for simulated monthly NEE at  $1 \times 1$  resolution from an example forward model. For each modeled monthly flux estimate (right), the grid-scale value was proportioned to the Forest Lands, Crop Lands and Other Lands sectors by weighting the flux according to the relative area of each land category (left) within a given grid cell. Model estimates at the level of the reporting zone were generated by then summing the flux across each sector within a given zone.



simulated fluxes to be summed by reporting zone and sector. Note that this approach meant that there could be discrepancies between how an inventory or model analysis might label the land surface and how we reported it (based on GLC-2000), but that compromise was necessary to accomplish the comparison.

## Results

### *Inventory estimates*

Overall, the data and methodology used herein for combining GHG inventory-based data on surface fluxes and carbon transfers across each sector and country suggest a  $-327 \text{ TgC yr}^{-1}$  (NEE) sink as the continental-scale carbon balance of North America over the 2000–2006 analysis period (Table 2). Our analysis finds that the continental-scale  $\text{CO}_2$  sink is driven primarily by  $\text{CO}_2$  uptake in the Forest Lands ( $-248 \text{ TgC yr}^{-1}$ ) and Crop Lands ( $-297 \text{ TgC yr}^{-1}$ ) sectors, with much of this sink offset by the source effect from the Other Lands sector ( $+218 \text{ TgC yr}^{-1}$ ). The large sink estimates for US forests ( $-244 \text{ TgC yr}^{-1}$ ) and croplands ( $-264 \text{ TgC yr}^{-1}$ ) include carbon removals in forest ( $-115 \text{ TgC yr}^{-1}$ ) and crop ( $-246 \text{ TgC yr}^{-1}$ ) harvested products, which are transferred to the Other Lands sector and contribute to a counteracting source ( $+207 \text{ TgC yr}^{-1}$ ). By comparison, the data show smaller sink estimates for forests ( $-31 \text{ TgC yr}^{-1}$ ) and croplands ( $-33 \text{ TgC yr}^{-1}$ ) in Canada, which are also offset in part by a source effect from the Other Lands sector ( $+20 \text{ TgC yr}^{-1}$ ). The sector-level NEE estimates for Mexico show a different pattern due to the flux estimates being primarily based on land use change effects. Here, Mexican forests are estimated as a net source to the atmosphere ( $+27 \text{ TgC yr}^{-1}$ ) whereas the data show a net sink effect from the Other Lands sector ( $-9.1 \text{ TgC yr}^{-1}$ ).

The detail on the inventory-based estimates of component fluxes that produce the patterns of NEE in the Forest Lands sector is illustrated in Fig. 3, and estimates for each reporting zone are provided in the Sup-


porting Information (Tables S1, Canada; S2, the US; and S3, Mexico). Forests in Canada and the US show carbon gains over the 2000–2006 time period in the dead organic matter pool ( $-40 \text{ TgC yr}^{-1}$  and  $-34 \text{ TgC yr}^{-1}$ , respectively) and the data suggest a large sink in live vegetation in US forests ( $-130 \text{ TgC yr}^{-1}$ ), but the inventory-based estimate of  $\Delta\text{Live}$  in Canada's managed forest area represents an overall decrease in carbon storage in the live vegetation pool ( $+47 \text{ TgC yr}^{-1}$ ). Harvest removals were  $-50 \text{ TgC yr}^{-1}$  for Canada and  $-115 \text{ TgC yr}^{-1}$  for the US. The forest sector of Mexico shows an overall loss of carbon over the time period of the inventory data (1993–2002) driven by biomass conversion ( $+18 \text{ TgC yr}^{-1}$ ) and soil carbon loss ( $+24 \text{ TgC yr}^{-1}$ ) from land use change, which is only partially offset by regenerating forests on abandoned agricultural lands ( $-2.7 \text{ TgC yr}^{-1}$ ) and net uptake by managed forests ( $-12 \text{ TgC yr}^{-1}$ ).

The inventory-based estimates of component crop NPP and harvest removals, along with  $\Delta\text{Soil}$  and NEE in the Crop Lands sector of each reporting zone in Canada and the US over the 2000–2006 time period are provided in the Supporting Information (Tables S4, Canada; and S5, the US). Overall, total carbon uptake by croplands (crop NPP) was more than six times greater in the US ( $-569 \text{ TgC yr}^{-1}$ ) than Canada ( $-89 \text{ TgC yr}^{-1}$ ). With small amounts of gain in cropland SOC stocks ( $\Delta\text{Soil}$ ) over this time period ( $-2.7 \text{ TgC yr}^{-1}$  for Canada and  $-18 \text{ TgC yr}^{-1}$  for the US), Crop Lands NEE was dominated by the crop harvest component ( $-30 \text{ TgC yr}^{-1}$  and  $-246 \text{ TgC yr}^{-1}$ , respectively). The concentration of the Crop Land NEE sink in the mid-continent region is illustrated in Fig. 3.

The magnitude of the contribution of forest and crop products to the national-/continental-scale net sink is a function of the relative amount of harvest that is stored over the time period, exported internationally, or returned to the atmosphere as non- $\text{CO}_2$  emissions. Most of the forest harvest contribution to the continental-scale sink (Table S6) is attributed to carbon storage in the US product pool ( $-39 \text{ TgC yr}^{-1}$ ) and the net export of forest harvest from Canada ( $-25 \text{ TgC yr}^{-1}$ ). On the fate of Canada and US harvested crop products, 79% is emitted as  $\text{CO}_2$  on the continent, with another 20% accounted for by international exports (a small amount is emitted as  $\text{CH}_4$  from livestock plus the contribution to stock increase in the human population). The contribution of harvested wood and crop products to the spatial pattern of NEE was assessed by calculating, for reporting zone, the net balance between product harvest and emissions (Fig. 4). This measure of each reporting zone's net product balance (NBP) highlights the large producers of forest (Northwest and Southeast) and crop (mid-West) products next to the

**Table 2** Inventory-based estimates of average annual total NEE ( $\text{TgC yr}^{-1}$ ) by country/sector, 2000–2006

Country	Sector			Total
	Forest lands	Crop lands	Other lands	
Canada	-31.00	-32.79	20.21	-43.58
US	-244.38	-264.32	206.69	-302.01
Mexico	27.47	<i>n/a</i>	-9.06	18.42
North America	-247.91	-297.11	217.84	-327.17




**Fig. 3** The spatial distribution of inventory-based estimates of average annual total NEE ( $\text{TgC yr}^{-1}$ ) across reporting zones, 2000–2006, for the (a) Forest Lands, (b) Crop Lands, and (c) Other Lands sectors, as well as for (d) all land area.

large consumers based on large human and livestock populations (California and Texas).

The magnitudes of the contribution of the various flux components to the total NEE from the Other Lands sector in each reporting zone over the 2000–2006 time period are illustrated in Fig. 3 and provided in the Supporting Information (Tables S7, Canada; and S8, the US). Grassland and settled areas contribute a small ‘background’ sink in Canada ( $-3.0 \text{ TgC yr}^{-1}$  and  $-3.1 \text{ TgC yr}^{-1}$ , respectively) and the US ( $-13 \text{ TgC yr}^{-1}$  and  $-27 \text{ TgC yr}^{-1}$ ). However, the emission of carbon that is transferred from the Forest Lands and Crop Lands sectors in the form of harvested (wood and crop) products overwhelm this small sink, resulting in a net  $\text{CO}_2$  source from the Other Lands sectors of both Canada and the US over this time period. Livestock emissions of  $\text{CO}_2$  related to the consumption of harvested

crop products account for the largest portion of this source in Canada ( $+20 \text{ TgC yr}^{-1}$ ) and the US ( $+181 \text{ TgC yr}^{-1}$ ). Most of the remaining Other Lands sector source effect is due to emissions from the decay of harvested wood products in Canada ( $+5.4 \text{ TgC yr}^{-1}$ ) and the US ( $+51 \text{ TgC yr}^{-1}$ ). A small amount of emissions is attributed to human consumption of harvested crop products in Canada ( $+1.8 \text{ TgC yr}^{-1}$ ) and the US ( $+15 \text{ TgC yr}^{-1}$ ). The magnitudes of the contribution of the various flux components to the total NEE from non-forest lands (Other Lands sector) in each reporting zone of Mexico over the 1993–2002 time period are provided in the Supporting Information (Table S9). The net sink effect estimated for the other lands sector of Mexico over this time period is driven by carbon storage in the soil pool ( $-16 \text{ TgC yr}^{-1}$ ) in agriculture, pasture, and forest plantation lands. Some of this sink is offset



**Fig. 4** The net product balance (NPB) between forest/crop product harvest and forest/crop product emissions ( $\text{TgC yr}^{-1}$ ), 2000–2006, for each reporting zone from the inventory-based estimates, shown for (a) forest harvest products balance,  $\text{NPB}_F = (H_R + H_E) + E_F$ , with croplands masked; (b) crop harvest products,  $\text{NPB}_C = H_R + (E_H + E_L)$ , with forest lands masked; and (c) all products,  $\text{NPB}_{\text{TOT}} = \text{NPB}_F + \text{NPB}_C$ . A negative value represents a net producing (exporting) zone and a positive value represents a net consuming (importing/emitting) zone.

by CO<sub>2</sub> emissions attributed to fuelwood harvest (+6.8 TgC yr<sup>-1</sup>), which is assumed here to be used within the same reporting zone that it was harvested.

#### Uncertainties and additional fluxes

Based on summing the upper and lower bounds on the range of uncertainty for each major component flux of the three sectors, the aggregate percent uncertainty on the inventory-based, continental-scale NEE estimate is approximately 77%, giving a range of -76 to -556 TgC yr<sup>-1</sup> (Table 3). At the sector-level, percent uncertainty on the inventory-based NEE estimates range from 17% for Crop Lands to 41% and 45% for Forest Lands and Other Lands, respectively. More detail on the uncertainty estimates for individual components, and the sources of these estimates, are given in the Supporting Information (Table S11). We also considered an additional -239 TgC yr<sup>-1</sup> NEE from 'best estimates' of additional components of the NA carbon budget that are not measured or estimated by the inventories, which are potentially significant but highly uncertain mostly due to the lack of available data. These estimates, primarily from those reported in the SOCCR (Pacala *et al.*, 2007), include additions to the continental-scale NEE of -120 TgC yr<sup>-1</sup> in woody encroachment in the US, -49 TgC yr<sup>-1</sup> for wetland ecosystems across NA, -25 TgC yr<sup>-1</sup> for sequestration in rivers and reservoirs of the US, and -45 TgC yr<sup>-1</sup> for DOC export from Canada and US rivers (Table S12). Given

**Table 3** The continental-scale, aggregate uncertainty around the inventory-based mean estimates of sector-level fluxes analyzed in this study, along with 'additional fluxes' not represented by the inventories. The detailed uncertainty estimates and additional fluxes for the various underlying components are provided in the Supporting Information (Tables S11 and S12)

Sector	Mean estimate	%	Uncertainty range relative to estimate	
			Lower bound (TgC yr <sup>-1</sup> )	Upper bound (TgC yr <sup>-1</sup> )
Forest lands	-245.30	41	-346.21	-144.40
Crop lands	-297.11	17	-347.54	-246.68
Other lands	217.84	45	120.82	314.86
Continental total	-324.57	77	-556.14	-76.21
Total 'additional fluxes'	-239.00	100	-572.93	0.00
Continental total w/'additional fluxes'	-563.57	86	-1050.93	-76.21

that each of these estimates carries at least 100% uncertainty, the aggregate additional flux could add anywhere from 0 to -573 TgC yr<sup>-1</sup> to our overall inventory-based estimate of continental-scale NEE.

#### Comparing inventory estimates to alternative scaling approaches

The mean model estimates (Table 4) from both the inverse (-931 TgC yr<sup>-1</sup> NEE) and forward (-511 TgC yr<sup>-1</sup>) approaches suggest a larger continental-scale total sink than does the result of our analysis of the various inventory-based data sets (-327 TgC yr<sup>-1</sup>, from Table 2). At the level of the reporting zone, different patterns among the three scaling approaches were compared by showing area-weighted NEE estimates for each sector in map format (Fig. 5). The range for mean annual NEE over North America among the inverse models was from a +15 TgC yr<sup>-1</sup> source to a -2190 TgC yr<sup>-1</sup> sink, with the five mid-range estimates clustering around a mean of -869 ± 223 TgC yr<sup>-1</sup>. The range of forward model estimates was from a small source (+29 TgC yr<sup>-1</sup>) to a large sink (-3210 TgC yr<sup>-1</sup>), with no real central tendency.

The mean modeled NEE estimates from the forward and inverse approaches (Table 4) follow a similar pattern of relative magnitude by country/sector as the inventory-based estimates, where the largest sink estimates are for the Forest Lands sector of the US (-282 TgC yr<sup>-1</sup> from the AIMs and -158 TgC yr<sup>-1</sup> from the TBMs), with smaller sink estimates for Canada's managed forest area (-151 TgC yr<sup>-1</sup> from the AIMs and -73 TgC yr<sup>-1</sup> from the TBMs). The mean NEE estimate for the Forest Lands sector of the US from the sets of AIMs represents a similar sink as we calculated from our analysis of the inventory data (-244 TgC yr<sup>-1</sup>), while the TBMs mean suggests a smaller sink. For Canada, both sets of models estimate a larger sink than the inventory-based results (-31 TgC yr<sup>-1</sup>) for the Forest Land sector. Both sets of models also estimate a smaller total Crop Lands sector sink for NA (-167 TgC yr<sup>-1</sup> from the AIMs and -134 TgC yr<sup>-1</sup> from the TBMs) than does the inventory-based approach (-295 TgC yr<sup>-1</sup>), which does not include in its estimate any data for the Crop Lands sector of Mexico. Compared to the relatively large CO<sub>2</sub> source from the Forest Lands sector of Mexico as estimated by the inventory data (+27 TgC yr<sup>-1</sup>), the mean Forest Lands sector NEE is near neutral (+0.9 TgC yr<sup>-1</sup>) from the AIMs and a small sink (-15 TgC yr<sup>-1</sup>) from the TBMs, although it should be noted that the time period covered by the inventory data (1993–2002) is different than that of the model estimates (~2000–2006). Beyond the Forest Lands and Crop

**Table 4** The count (n), mean and standard deviation (SD) of average annual NEE estimates (TgC yr<sup>-1</sup>), 2000–2006 by country and sector, for the sets of inverse and forward models. The mean estimates from the inventory-based approach (from Table 2) for each country and sector are included for comparison

Country/Sector	Inverse models			Forward models			Inventory-based
	<i>n</i>	Mean	SD	<i>n</i>	Mean	SD	Mean estimate
Canada total	7	-237.6	96.7	15	-124.6	205.5	-43.6
Forestland	7	-150.9	55.4	15	-73.3	141.3	-31.0
Cropland	7	-35.5	24.3	15	-22.1	27.5	-32.8
Other	7	-51.2	28.3	15	-29.3	41.0	20.2
U. S. Total	7	-685.1	573.7	17	-357.0	575.5	-302.0
Forestland	7	-282.0	214.1	17	-157.6	309.5	-244.4
Cropland	7	-136.8	124.0	17	-94.6	160.3	-264.3
Other	7	-266.2	263.2	17	-104.8	127.9	206.7
Mexico total	7	-8.7	159.2	12	-29.0	71.8	18.4
Forestland	7	0.9	63.6	12	-15.1	48.1	27.5
Cropland	7	5.5	33.2	12	-17.5	33.0	<i>n/a</i>
Other	7	-15.1	63.8	12	3.6	34.1	-9.1
N. America total	7	-931.3	670.3	12	-510.7	729.3	-327.2
Forestland	7	-432.1	254.1	12	-246.0	419.2	-247.9
Cropland	7	-166.8	150.9	12	-134.2	194.3	-297.1
Other	7	-332.5	301.3	12	-130.5	151.8	217.8

Lands sector comparisons, it is primarily the difference in NEE estimates for the Other Lands sector that is responsible for the larger continental-scale sink estimates from the model means vs. the inventory-based data. At the continental-scale, the model mean NEE estimates from the AIMs (-333 TgC yr<sup>-1</sup>) and TBMs (-131 TgC yr<sup>-1</sup>) show a large sink in the Other Lands sector, whereas the results of the inventory-based methodology used herein suggests a large source (+218 TgC yr<sup>-1</sup>).

## Discussion

### *Inventory-based estimates*

Our GHG inventory-based results are derived from, and so are generally consistent with, recent inventory-based updates of the carbon budgets reported for Canada forests (Pan *et al.*, 2011; Stinson *et al.*, 2011), US forests (Heath *et al.*, 2011; Pan *et al.*, 2011) and agriculture (West *et al.*, 2011), and the agriculture and forest sector in Mexico (deJong *et al.*, 2010). The new information provided in this study comes from the combination of those national- and sector- specific estimates into a continental-scale analysis, while using a novel conceptual model to estimate land-atmosphere exchange of CO<sub>2</sub> at the sub-national scale. As a result, the inventory-based data and the methodology used in this study suggest considerable spatial variability

in NEE estimates across sectors and reporting zones (Fig. 3). The spatial patterns are driven both by the estimated direct, vertical surface fluxes as well as the lateral transfer of carbon between sectors in the form of harvested products (Fig. 4). The spatial patterns show a negative balance (i.e., sink effect) between product emissions and harvest in reporting zones that have relatively smaller human and livestock populations but productive forests and croplands with high harvest rates (and vice versa).

The largest Forest Lands sector CO<sub>2</sub> sinks are located primarily on the west coast and in the south-east of the US, and these estimates are similar in magnitude to sub-regional analyses by Turner *et al.* (2011) and Masek & Collatz (2006). Despite covering roughly similar area, Canada shows a much smaller magnitude sink in the Forest Lands sector than does the US. Although some of this difference could be related to methodology (Kurz *et al.*, 2009; Heath *et al.*, 2011), Canada's forests are likely to be storing less carbon than US forests due to older age class structure, lower growth rates and higher frequency and severity of disturbances in boreal forests vs. temperate forests (Kurz *et al.*, 2008; Stinson *et al.*, 2011). All the reporting zones for Mexico show a small source from the forest sector, with the largest sources in southern states that have higher proportions of lowland tropical forest, where most of the forest clearing has occurred (deJong *et al.*, 2010). The analysis of the net




Fig. 5 Mean area-weighted average annual NEE ( $\text{g C m}^{-2} \text{yr}^{-1}$ ), 2000–2006 for the Forest Lands, Crop Lands and Other Lands sectors, along with all land (total), in each reporting zone, from inventory-based estimates against mean results from the sets of terrestrial biosphere (forward) models and inverse models.

land use change impact implies that, at the national-level, emissions from biomass conversion across Mexico are outpacing uptake from forests re-growing after agricultural abandonment.

The continental-scale mapping of NEE for the Crop Lands sector reflects the pattern of strong net carbon uptake over the mid-western US, as discussed in other studies (Corbin *et al.*, 2010; West *et al.*, 2010). Although we assign this uptake to the Crop Lands sector sink, most (79%) of this carbon is returned to the atmosphere after consumption and respiration by humans and livestock (West *et al.*, 2009) within North America, which we attribute to the Other Lands sector source. Nearly all the remaining balance of harvested crop product C is exported internationally. Although emissions of this remaining balance are not counted from the atmospheric perspective over North America, these emissions will occur in other countries. Thus, from a global atmospheric perspective, the net contribution of harvested crop product C to NEE is near neutral.

#### Comparison to model estimates

The mean model estimates from both the forward (TBMs) and inverse (AIMs) approaches suggest a much stronger overall NA sink than the inventory-based estimate. Yet model estimates generally do follow similar spatial patterns as the inventory-based data where the strongest sinks are found in US forests on the east and west coasts and in croplands of the mid continent, with a smaller source from the tropical area of southern Mexico (Fig. 5). However, the model vs. inventory differences are mostly in the magnitude of the estimates, where the sector-specific model means suggest (1) a larger sink over forested regions, (2) a smaller sink over crop land areas, and (3) a substantial contribution of non-forest/non-cropland areas to the continental-scale sink (Table 4).

At the national-level, the breakdown of model means for the Forest Lands sectors show good agreement with the inventory-based estimate for the US, but a much

larger sink than that estimated by inventory-based modeling for the Forest Lands sector in Canada. Inventory-based studies indicate that CO<sub>2</sub> uptake in Canada's forests is being increasingly offset by emissions due to disturbance (Kurz & Apps, 1999; Kurz *et al.*, 2008; Stinson *et al.*, 2011), but our comparisons here suggest that the impacts of these disturbances are not being resolved by the model approaches. In contrast to the Forest Lands sector comparison, the model means estimate less than half of the sink strength compared to the inventory-based estimate for the US Crop Lands sector. There is clearly information in the CO<sub>2</sub> observations indicating a strong drawdown in the crop intensive region of the US (Corbin *et al.*, 2010; Crevoisier *et al.*, 2010), but the model ensembles analyzed herein appear to be underestimating its strength, relative to the inventory estimates.

The difference in the sign and magnitude between the inventory and model approaches in the case of the Other Lands sector highlights (1) our inventory-based approach for allocating product respiration and decomposition based on populations of humans and livestock and (2) the data gaps and uncertainties associated with GHG inventory-based estimates of carbon stocks and fluxes outside of managed forest and agricultural lands. Although a subset of the TBMs included herein considers forest and/or crop product emissions, none considers the lateral transfer of these products (i.e. product emissions occur in the same grid cell as growth and harvest). AIMS derive the 'land flux' after prescribing the fossil fuel and fire emissions. In practice, the land flux thus includes the product sources. However, it is generally acknowledged that uncertainty remains high for inversion-based flux estimates at the sub-continental scale (Butler *et al.*, 2010; Bruhwiler *et al.*, 2011). As such, source areas associated with the respiration of harvested products may not be spatially resolved. On the other hand, potential sinks in the Other Lands sector that may be included in the model estimates could be missing or are of highly uncertain magnitude based on GHG inventory methods. For example, the SOCCR reports an additional 120 TgC yr<sup>-1</sup> of uptake through woody encroachment in the US, but other field-based studies (Goodale & Davidson, 2002b; Jackson *et al.*, 2002) do not support a sink of that magnitude. Further, it is not clear how much of this mechanism is captured in the inventory sampling if and where it is occurring. It is evident in the US forest statistics that a large proportion of the increase in US forest land has occurred in the West. Due to long re-measurement periods and changes in methods over recent time periods, however, it is not possible to determine how much of that increase is directly attributable to woody encroachment.

### Synthesis

Multi-method flux comparisons over other large regions are similar to our comparison in several respects. In both Europe (Janssens *et al.*, 2003) and China (Piao *et al.*, 2009), the land base was a sink for carbon and represented a significant proportion of fossil fuel emissions (7–12% in Europe and 28–37% in China). In both cases the inversion-based sink estimate was about double the inventory or process model-based sink estimates. An updated, multi-sector study of the European C balance (Schulze *et al.*, 2010), based primarily on inventory methods, suggests that C sinks (e.g., forests and grassland) are largely offset by emissions (e.g., from croplands). As with our North American study, the lateral movement of harvested products was also considered to be a large influence on the spatial distribution of sources and sinks in Europe (Ciais *et al.*, 2006; Luyssaert *et al.*, 2010). Over the 2000–2006 time period, our national-level inventory-based NEE estimates represent approximately 29% and 19% of fossil fuel emissions for Canada (0.15 PgC yr<sup>-1</sup> ± 4%) and the US (1.56 PgC yr<sup>-1</sup> ± 4%), respectively (Boden *et al.*, 2010). Our inventory-based NEE estimate for Mexico adds approximately 18% to the fossil fuel source from that country (0.11 PgC yr<sup>-1</sup> ± 4%). Including the 'best estimates' for additional component fluxes not measured in the inventories would increase the inventory-based sink estimate to approximately 31% of total continental-scale fossil emissions (1.83 PgC yr<sup>-1</sup>). Mean NEE estimates from the ensembles of TBMs and AIMS represent land-based sinks that offset 28% and 51%, respectively, of total continental-scale fossil emissions (1.83 PgC yr<sup>-1</sup>).

A large land-based CO<sub>2</sub> sink over NA has been a persistent feature of inversion analyses and comparisons of inversions to bottom-up estimates at the regional (Hayes *et al.*, 2011; Turner *et al.*, 2011) and continental (Pacala *et al.*, 2001) scales have suggested that it is an overestimate. First, the biases in vertical mixing in the transport models could lead to the overestimates of the source strength in tropical latitudes and overestimates of the sink strength in mid latitudes (Stephens *et al.*, 2007; Gatti *et al.*, 2010). Second, overestimates of NA west coast boundary conditions for CO<sub>2</sub> concentration may force the AIMS to create an artificial sink to maintain consistency with the measured CO<sub>2</sub> observations encountered further east (Göckede *et al.*, 2010; Schuh *et al.*, 2010). With respect to the forward modeling approach, the extremely large range in the flux estimates from the TBMs can be attributed to variation in model formulation and process representation along with differences in the climate and land use data sets used as model drivers (Schwalm *et al.*, 2010; Huntzinger

ger *et al.*, in press). In many cases, the large estimated sinks in TBMs are associated with assumptions of robust favorable effects of rising CO<sub>2</sub> on vegetation growth, but the magnitude of the effect of this mechanism remains highly uncertain (Joos *et al.*, 2002; Girardin *et al.*, 2011). The relative impact of any CO<sub>2</sub> fertilization effect is generally not possible to ascertain from the inventory data. In the Canada forest inventory approach, the species and site specific yield curves used to model NPP would not likely capture this effect. The US forest inventory should, in theory, capture this effect between re-measurement periods, but it is impossible to separate it from all other effects on growth.

This study's inventory-based, continental-scale NEE estimate of  $-327 \text{ TgC yr}^{-1}$  for the early 21st Century is generally lower than estimates from previous decades, which range from  $-350$  to  $-750 \text{ TgC yr}^{-1}$  (Houghton *et al.*, 1999; Pacala *et al.*, 2001, 2007; Goodale *et al.*, 2002a). The SOCCR is the most recent and comprehensive study, which yielded a NEE estimate of  $-500 \pm 250 \text{ TgC yr}^{-1}$  for NA in ca. 2003 (Pacala *et al.*, 2007). Although the sector-level NEE estimates presented herein are generally consistent with those reported for 'forests' and 'agricultural soils' in the SOCCR, the largest difference contributing to the lower continental carbon sink estimate here is that we did not include the large but highly uncertain additional fluxes associated with land-based sinks of atmospheric CO<sub>2</sub> (Table 3).

We would need to assume a large contribution of these non-inventoried 'additional fluxes' on top of the inventory-based sink estimate to approach the magnitude suggested by the means of the model ensembles analyzed in this study. For example, adding the 'best guess' of these non-inventoried 'additional fluxes' gives

an estimate of NEE ( $-564 \text{ TgC yr}^{-1}$ ; Table 3) that is similar to the mean of the TBMs ensemble ( $-511 \text{ TgC yr}^{-1}$ ; Table 4). The mean NEE estimate of the AIMs ensemble ( $-931 \text{ TgC yr}^{-1}$ ; Table 4) is found only near the extreme lower bound of the uncertainty around the inventory-based NEE estimate for the 'continental total w/'additional fluxes' ( $-1051 \text{ TgC yr}^{-1}$ ; Table 3). However, given that this analysis highlights the (1) uncertainties in component fluxes, (2) mismatches in spatial patterns, and (3) large spread in estimates across models, any convergence between the approaches would not necessarily occur for the 'right' reasons. Rather, this study draws attention to those components of the NA carbon budget that require more careful study through measurement and inventory methods. Regarding the modeling approaches, the comparisons here strongly suggest the need to better understand the causes underlying the large spread in estimates, most likely achieved through formal and controlled (i.e. common protocol) model inter-comparison studies informed by benchmarking frameworks based on reliable measurements and observational data sets.

This study highlights the differences in three general scaling approaches to NEE (inventory, forward and inverse modeling), and by comparing and evaluating their estimates several strengths and weaknesses emerge (Table 5). Our study suggests that, even considering the data gaps and uncertainties, the inventory-based approach to estimating NEE can still provide a substantial amount of important information at the sub-continental scale, and help inform estimates of both vertical and lateral transfers of most key carbon budget components. The strength of the inventory-based measurement approach is primarily its reliance on a large

**Table 5** A comparison of the strengths and weaknesses of alternative NEE scaling approaches (inventory-based, AIMs and TBMs)

	Inventory-based	Atmospheric inversion models (AIMs)	Terrestrial biosphere models (TBMs)
Strengths	<ol style="list-style-type: none"> <li>1) Employs a large number of repeated biomass measurements</li> <li>2) Allows estimation of product-related C sources</li> </ol>	<ol style="list-style-type: none"> <li>1) assimilates measurements of atmospheric CO<sub>2</sub> concentration</li> <li>2) Employs atmospheric mass balance</li> </ol>	<ol style="list-style-type: none"> <li>1) Processes are represented so attribution is possible</li> <li>2) Sensitive to interannual variation in climate</li> <li>3) Many opportunities for validation</li> </ol>
Weaknesses	<ol style="list-style-type: none"> <li>1) Not all C pools are measured</li> <li>2) Possible undersampling</li> <li>3) Limited attribution ability</li> <li>4) Missing NEE of unmanaged ecosystems</li> <li>5) Poorly resolved temporally</li> </ol>	<ol style="list-style-type: none"> <li>1) Transport model uncertainty</li> <li>2) Limited number of CO<sub>2</sub> measurements</li> <li>3) Low spatial resolution</li> <li>4) Limited attribution ability</li> </ol>	<ol style="list-style-type: none"> <li>1) Many inputs, each with their own uncertainty</li> <li>2) Many parameters, each with their own uncertainty</li> <li>3) Spatial resolution may not resolve management scale disturbances</li> </ol>

number of ground-based measurements of components useful to estimate carbon stocks and stock changes. Although there are benefits in retaining independence among approaches for estimating carbon fluxes, progress can also be made by more formally integrating them. For example, TBMs are increasingly making use of inventory and remote sensing data for model drivers, parameterization, calibration, and validation (e.g. Hurtt *et al.*, 2002; Running *et al.*, 2004). Such integrated 'bottom up' modeling frameworks could provide the initial land surface flux estimates for inversion analyses and, in turn, information about errors in predicted CO<sub>2</sub> concentration would inform further model development. Furthermore, observations and inventory-based measurements can provide critical benchmarking data sets for model evaluation (Randerson *et al.*, 2009). Ultimately, confidence in our ability to understand and predict the role of the NA carbon cycle in the global climate system will increase as the estimates from these different approaches begin to more closely converge and are combined in more fully integrated modeling systems.

### Acknowledgements

Research was conducted in part at Oak Ridge National Laboratory, and supported by the US Department of Energy (DOE), Office of Science, Biological and Environmental Research. Oak Ridge National Laboratory is managed by UT-Battelle for DOE under contract DE-AC05-00OR22725. The research reported in this paper was supported by multiple sources, including USDA CSREES grant 2008-35615-18959, NASA New Investigator Program grant NNX10AT66G and NASA Terrestrial Ecology Program grant NNX09AL51G. The authors would like to thank all of the modeling teams participating in the North American Carbon program and providing simulation result for this analysis through the Regional – Continental Interim Synthesis activity (<http://nacarbon.org>) and Robert Andres of the Los Alamos National Laboratory for provision of the data on fossil fuel emissions. We also acknowledge the efforts of Chris Williams, Jim Collatz, and the anonymous reviewers for greatly improving the quality of this manuscript through their added insight and constructive criticism.

### References

Bachelet D, Neilson RP, Hickler T *et al.* (2003) Simulating past and future dynamics of natural ecosystems in the United States. *Global Biogeochem. Cycles*, **17**, 1045, 21 pp, doi: 10.1029/2001GB001508.

Baker DF, Law RM, Gurney KR *et al.* (2006) TransCom 3 inversion intercomparison: impact of transport model errors on the interannual variability of regional CO<sub>2</sub> fluxes, 1988–2003. *Global Biogeochemical Cycles*, **20**, GB1002, 17 pp.

Baker IT, Prihodko L, Denning AS, Goulden M, Miller S, da Rocha HR (2008) Seasonal drought stress in the Amazon: Reconciling models and observations. *J. Geophys. Res.*, **113**, G00B01, 10 pp, doi: 10.1029/2007JG000644.

Bartholome E, Belward AS (2005) GLC2000: a new approach to global land cover mapping from earth observation data. *International Journal of Remote Sensing*, **26**, 1959–1977.

Bechtold WA, Patterson PL (2005) *The Enhanced Forest Inventory and Analysis Program – National Sampling Design and Estimation Procedures*. SRS GTR-80. USDA Forest Service, Southern Research Station, Asheville, NC, USA.

Boden TA, Marland G, Andres RJ (2010) *Global, Regional, and National Fossil-Fuel CO<sub>2</sub> Emissions*. Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, U.S. Department of Energy, Oak Ridge, TN, USA, doi: 10.3334/CDIAC/00001\_V2010.

Bondeau A, Smith PC, Zaehle S *et al.* (2007) Modelling the role of agriculture for the 20th century global terrestrial carbon balance. *Global Change Biology*, **13**, 679–706.

Bruhwieler LM, Michalak AM, Tans PP (2011) Spatial and temporal resolution of carbon flux estimates for 1983–2002. *Biogeosciences*, **8**, 1309–1331.

Butler MP, Davis KJ, Denning AS, Kawa SR (2010) Using continental observations in global atmospheric inversions of CO<sub>2</sub>: North American carbon sources and sinks. *Tellus Series B-Chemical and Physical Meteorology*, **62**, 550–572.

Chapin FS III, Woodwell GM, Randerson JT *et al.* (2006) Reconciling carbon-cycle concepts, terminology, and methods. *Ecosystems*, **9**, 1041–1050.

Chevallier F, Bréon F-M, Rayner PJ (2007) Contribution of the Orbiting Carbon Observatory to the estimation of CO<sub>2</sub> sources and sinks: Theoretical study in a variational data assimilation framework. *J. Geophys. Res.*, **112**, D09307, 11 pp, doi: 10.1029/2006JD007375.

Ciais P, Canadell JG, Luysaert S *et al.* (2010) Can we reconcile atmospheric estimates of the Northern terrestrial carbon sink with land-based accounting? *Current Opinion in Environmental Sustainability*, **2**, 225–230.

Ciais P, Borges AV, Abril G, Meybeck M, Folberth G, Hauglustaine D, Janssens IA (2006) The impact of lateral carbon fluxes on the European carbon balance. *Biogeosciences Discuss*, **3**, 1529–1559.

Corbin KD, Denning AS, Lokupitiya EY *et al.* (2010) Assessing the impact of crops on regional CO<sub>2</sub> fluxes and atmospheric concentrations. *Tellus Series B-Chemical and Physical Meteorology*, **62**, 521–532.

Crevoisier C, Sweeney C, Gloor M, Sarmiento JL, Tans PP (2010) Regional US carbon sinks from three-dimensional atmospheric CO<sub>2</sub> sampling. *Proceedings of the National Academy of Sciences*, **107**, 18348–18353.

deJong B, Anaya C, Masera O *et al.* (2010) Greenhouse gas emissions between 1993 and 2002 from land-use change and forestry in Mexico. *Forest Ecology and Management*, **260**, 1689–1701.

Deng F, Chen JM, Ishizawa M *et al.* (2007) Global monthly CO<sub>2</sub> flux inversion with a focus over North America. *Tellus Series B-Chemical and Physical Meteorology*, **59**, 179–190.

Denning AS, Oren R, McGuire AD *et al.* (2005) *Science implementation strategy for the North American carbon program*. Report of the NACP Implementation Strategy Group of the U.S. Carbon Cycle Interagency Working Group, U.S. Carbon Cycle Science Program, Washington, DC.

Eggelston HS, Buendia L, Miwa K, Ngara T, Tanabe K. (2006) *Guidelines for National Greenhouse Gas Inventories*. IPCC National Greenhouse Gas Inventories Programme, Hayama, Japan.

EPA (2011) Inventory of U.S. Greenhouse Gas Emissions and Sinks: 1990–2009. USEPA #430-R-11-005. U.S. Environmental Protection Agency, Washington, DC. Available at: <http://www.epa.gov/climatechange/emissions/usinventoryreport.html> (accessed 15 April 2011).

Environment Canada (2011) National Inventory Report 1990–2009: Greenhouse Gas Sources and Sinks in Canada. The Government of Canada's Submission to the UN Framework Convention on Climate Change. Environment Canada, Ottawa, ON. Available at: <http://www.ec.gc.ca/ges-ghg/> (accessed 16 May 2011).

Fan S, Gloor M, Mahlman J, Pacala S, Sarmiento J, Takahashi T, Tans P (1998) A large terrestrial carbon sink in North America implied by atmospheric and oceanic carbon dioxide data and models. *Science*, **282**, 442–446.

Gatti LV, Miller JB, D'Amelio MTS *et al.* (2010) Vertical profiles of CO<sub>2</sub> above eastern Amazonia suggest a net carbon flux to the atmosphere and balanced biosphere between 2000 and 2009. *Tellus Series B-Chemical and Physical Meteorology*, **62**, 581–594.

Girardin MP, Bernier PY, Raulier F, Tardif JC, Conciatori F, Guo XJ (2011) Testing for a CO<sub>2</sub> fertilization effect on growth of Canadian boreal forests. *Journal of Geophysical Research-Biogeosciences*, **116**, G01012, 16 pp, doi: 10.1029/2010JG001287.

Göckede M, Turner DP, Michalak AM, Vickers D, Law BE (2010) Sensitivity of a subregional scale atmospheric inverse CO<sub>2</sub> modeling framework to boundary conditions. *Journal of Geophysical Research*, **115**, D24112.

Goodale CL, Apps MJ, Birdsey RA *et al.* (2002a) Forest carbon sinks in the Northern Hemisphere. *Ecological Applications*, **12**, 891–899.

Goodale CL, Davidson EA (2002b) Carbon cycle: uncertain sinks in the shrubs. *Nature*, **418**, 593–594.

Gurney KR, Law RM, Denning AS *et al.* (2002) Towards robust regional estimates of CO<sub>2</sub> sources and sinks using atmospheric transport models. *Nature*, **415**, 626–630.



- Gurney KR, Law RM, Denning AS *et al.* (2004) Transcom 3 inversion intercomparison: model mean results for the estimation of seasonal carbon sources and sinks. *Global Biogeochemical Cycles*, **18**, GB1010, 18 pp, doi: 10.1029/2003GB002111.
- Hayes DJ, McGuire AD, Kicklighter DW, Gurney KR, Burnside TJ, Melillo JM (2011) Is the northern high latitude land-based CO<sub>2</sub> sink weakening? *Global Biogeochemical Cycles*, **25**, GB3018, 14 pp, doi: 10.1029/2010GB003813.
- Heath LS, Smith JE, Skog KE, Nowak DJ, Woodall CW (2011) Managed forest carbon estimates for the US greenhouse gas inventory, 1990–2008. *Journal of Forestry*, **109**, 167–173.
- Houghton RA, Hackler JL, Lawrence KT (1999) The US carbon budget: contributions from land-use change. *Science*, **285**, 574–578.
- Huntzinger DN, Post WM, Wei Y *et al.* (in press) North american carbon project (NACP) regional interim synthesis: terrestrial biospheric model intercomparison. *Ecological Modeling*.
- Hurrut GC, Pacala SW, Moorcroft PR, Caspersen J, Shevliakova E, Houghton RA, Moore B 3rd (2002) Projecting the future of the U.S. carbon sink. *Proceedings of the National Academy of Sciences of the United States of America*, **99**, 1389–1394.
- Jackson RB, Banner JL, Jobbagy EG, Pockman WT, Wall DH (2002) Ecosystem carbon loss with woody plant invasion of grasslands. *Nature*, **418**, 623–626.
- Janssens IA, Freibauer A, Ciais P *et al.* (2003) Europe's terrestrial biosphere absorbs 7 to 12% of European anthropogenic CO<sub>2</sub> emissions. *Science*, **300**, 1538–1542.
- Joos F, Prentice IC, House JI (2002) Growth enhancement due to global atmospheric change as predicted by terrestrial ecosystem models: consistent with US forest inventory data. *Global Change Biology*, **8**, 299–303.
- Ju W, Chen JM, Black TA, Barr AG, Liu J, Chen B (2006) Modelling multi-year coupled carbon and water fluxes in a boreal aspen forest. *Agricultural and Forest Meteorology*, **140**, 136–151.
- King AW, Dilling L, Zimmerman GP *et al.* (2007). What is the carbon cycle and why care? In: *The First State of the Carbon Cycle Report (SOCCR): The North American Carbon Budget and Implications for the Global Carbon Cycle*. A Report by the U.S. Climate Change Science Program and the Subcommittee on Global Change Research (eds King AW, Dilling L, Zimmerman GP, Fairman DM, Houghton RA, Marland G, Rose AZ, Wilbanks TJ). pp. 15–20. National Oceanic and Atmospheric Administration, National Climatic Data Center, Asheville, NC, USA.
- Krinner G, Viovy N, de Noblet-Ducoudré N *et al.* (2005) A dynamic global vegetation model for studies of the coupled atmosphere-biosphere system. *Global Biogeochemical Cycles*, **19**, GB1015, 33 pp, doi: 10.1029/2003GB002199.
- Kucharik CJ, Foley JA, Delire C *et al.* (2000) Testing the performance of a dynamic global ecosystem model: Water balance, carbon balance, and vegetation structure. *Global Biogeochemical Cycles*, **14**, 795–825.
- Kurz WA, Apps MJ (1999) A 70-year retrospective analysis of carbon fluxes in the Canadian forest sector. *Ecological Applications*, **9**, 526–547.
- Kurz WA, Dymond CC, Stinson G *et al.* (2008) Mountain pine beetle and forest carbon feedback to climate change. *Nature*, **452**, 987–990.
- Kurz WA, Dymond CC, White TM *et al.* (2009) CBM-CFS3: a model of carbon-dynamics in forestry and land-use change implementing IPCC standards. *Ecological Modelling*, **220**, 480–504.
- Lokupitiya RS, Zupanski D, Denning AS, Kawa SR, Gurney KR, Zupanski M (2008) Estimation of global CO<sub>2</sub> fluxes at regional scale using the maximum likelihood ensemble filter. *Journal of Geophysical Research-Atmospheres*, **113**, D20110, 19 pp, doi: 10.1029/2007JD009679.
- Luyssaert S, Ciais P, Piao SL *et al.* (2010) The European carbon balance. Part 3: forests. *Global Change Biology*, **16**, 1429–1450.
- Masek JG, Collatz GJ (2006) Estimating forest carbon fluxes in a disturbed southeastern landscape: integration of remote sensing, forest inventory, and biogeochemical modeling. *Journal of Geophysical Research-Biogeosciences*, **111**, G01006, 15 pp, doi: 10.1029/2005JG000062.
- Michalak AM, Bruhwiler L, Tans PP (2004) A geostatistical approach to surface flux estimation of atmospheric trace gases. *J. Geophys. Res.*, **109**, D14109, 19 pp, doi: 10.1029/2003JD004422.
- Myneni RB, Dong J, Tucker CJ *et al.* (2001) A large carbon sink in the woody biomass of Northern forests. *Proceedings of the National Academy of Sciences of the United States of America*, **98**, 14784–14789.
- Pacala S, Birdsey RA, Bridgman SD *et al.* (2007) The North American carbon budget past and present. In: *The First State of the Carbon Cycle Report (SOCCR): The North American Carbon Budget and Implications for the Global Carbon Cycle* (eds King AW, Dilling L, Zimmerman GP, Fairman DM, Houghton RA, Marland G, Rose AZ, Wilbanks TJ). pp. 29–36. National Oceanic and Atmospheric Administration, National Climatic Data Center, Asheville, NC, USA.
- Pacala SW, Hurrut GC, Baker D *et al.* (2001) Consistent land- and atmosphere-based US carbon sink estimates. *Science*, **292**, 2316–2320.
- Pan Y, Birdsey RA, Fang J *et al.* (2011) A large and persistent carbon sink in the world's forests. *Science*, **333**, 988–993.
- Parson EA, Haas PM, Levy MA (1992) A summary of the major documents signed at the Earth Summit and the Global Forum. *Environment*, **34**, 12–15.
- Peters W, Jacobson AR, Sweeney C *et al.* (2007) An atmospheric perspective on North American carbon dioxide exchange: carbonTracker. *Proceedings of the National Academy of Sciences of the United States of America*, **104**, 18925–18930.
- Peylin P, Bousquet P, Le Quééré C *et al.* (2005) Multiple constraints on regional CO<sub>2</sub> flux variations over land and oceans. *Global Biogeochemical Cycles*, **19**, GB1011, 21 pp, doi: 10.1029/2003GB002214.
- Piao S, Fang J, Ciais P, Peylin P, Huang Y, Sitch S, Wang T (2009) The carbon balance of terrestrial ecosystems in China. *Nature*, **458**, 1009–1013.
- Randerson JT, Hoffman FM, Thornton PE *et al.* (2009) Systematic assessment of terrestrial biogeochemistry in coupled climate – carbon models. *Global Change Biology*, **15**, 2462–2484.
- Randerson JT, Thompson MV, Conway TJ, Fung IY, Field CB (1997) The contribution of terrestrial sources and sinks to trends in the seasonal cycle of atmospheric carbon dioxide. *Global Biogeochem. Cycles*, **11**, 535–560.
- Reichstein M, Falge E, Baldocchi D *et al.* (2005) On the separation of net ecosystem exchange into assimilation and ecosystem respiration: review and improved algorithm. *Global Change Biology*, **11**, 1424–1439.
- Rodenbeck C, Houweling S, Gloor M, Heimann M (2003) CO<sub>2</sub> flux history 1982–2001 inferred from atmospheric data using a global inversion of atmospheric transport. *Atmospheric Chemistry and Physics*, **3**, 1919–1964.
- Running SW, Nemani RR, Heinsch FA, Zhao M, Reeves M, Hashimoto H (2004) A continuous satellite-derived measure of global terrestrial primary production. *BioScience*, **54**, 547–560.
- Schuh AE, Denning AS, Corbin KD *et al.* (2010) A regional high-resolution carbon flux inversion of North America for 2004. *Biogeosciences*, **7**, 1625–1644.
- Schulze ED, Ciais P, Luyssaert S *et al.* (2010) The European carbon balance. Part 4: integration of carbon and other trace-gas fluxes. *Global Change Biology*, **16**, 1451–1469.
- Schwalm CR, Williams CA, Schaefer K *et al.* (2010) A model-data intercomparison of CO<sub>2</sub> exchange across North America: results from the North American carbon program site synthesis. *Journal of Geophysical Research*, **115**, G00H05, 22 pp, doi: 10.1029/2009JG001229.
- Stephens BB, Gurney KR, Tans PP *et al.* (2007) Weak northern and strong tropical land carbon uptake from vertical profiles of atmospheric CO<sub>2</sub>. *Science*, **316**, 1732–1735.
- Stinson G, Kurz WA, Smyth CE *et al.* (2011) An inventory-based analysis of Canada's managed forest carbon dynamics, 1990 to 2008. *Global Change Biology*, **17**, 2227–2244.
- Smith JE, Heath LS, Nichols MC (2010) *U.S. forest carbon calculation tool: forestland carbon stocks and net annual stock change*. Revised for use with FIADB4.0. NRS GTR-13. USDA Forest Service, Northern Research Station, Newtown Square, PA, USA. [DVD-ROM].
- Smith JE, Heath LS, Skog KE, Birdsey RA (2006) *Methods for calculating forest ecosystem and harvested carbon with standard estimates for forest types of the United States*. Gen. Tech. Rep. NE-343. U.S. Department of Agriculture, Forest Service, Northeastern Research Station, Newtown Square, PA.
- Smith WB, Miles PD, Perry CH, Pugh SA (2009) *Forest resources of the United States, 2007*. WO GTR-78. USDA Forest Service, Washington Office, Washington, DC, USA.
- Thornton PE, Doney SC, Lindsay K *et al.* (2009) Carbon-nitrogen interactions regulate climate-carbon cycle feedbacks: results from an atmosphere-ocean general circulation model. *Biogeosciences*, **6**, 2099–2120.
- Tian H, Melillo J, Lu C *et al.* (2011) China's terrestrial carbon balance: Contributions from multiple global change factors. *Global Biogeochem. Cycles*, **25**, GB1007, 16 pp, doi: 10.1029/2010GB003838.
- Turner DP, GÖckede M, Law BE *et al.* (2011) Multiple constraint analysis of regional land – surface carbon flux. *Tellus Series B-Chemical and Physical Meteorology*, **63**, 207–221.
- van der Werf GR, Randerson JT, Giglio L, Collatz GJ, Kasibhatla PS, Arellano AF Jr (2006) Interannual variability in global biomass burning emissions from 1997 to 2004. *Atmos. Chem. Phys.*, **6**, 3423–3441.
- West T, Marland G, Singh N, Bhaduri B, Roddy A (2009) The human carbon budget: an estimate of the spatial distribution of metabolic carbon consumption and release in the United States. *Biogeochemistry*, **94**, 29–41.

- West TO, Bandaru V, Brandt CC, Schuh AE, Ogle SM (2011) Regional uptake and release of crop carbon in the United States. *Biogeosciences Discuss*, **8**, 631–654.
- West TO, Brandt CC, Baskaran LM *et al.* (2010) Cropland carbon fluxes in the United States: increasing geospatial resolution of inventory-based carbon accounting. *Ecological Applications*, **20**, 1074–1086.
- Wofsy SC, Harriss RC (2002) The North American carbon program (NACP). Report of the NACP Committee of the U.S. Interagency Carbon Cycle Science Program. US Global Change Research Program, Washington, DC
- Xiao J, Zhuang Q, Baldocchi DD *et al.* (2008) Estimation of net ecosystem carbon exchange for the conterminous United States by combining MODIS and AmeriFlux data. *Agricultural and Forest Meteorology*, **148**, 1827–1847.
- Yang X, Wittig V, Jain AK, Post W (2009) Integration of nitrogen cycle dynamics into the Integrated Science Assessment Model for the study of terrestrial ecosystem responses to global change. *Global Biogeochem. Cycles*, **23**, GB4029, 18 pp. doi: 10.1029/2009GB003474.
- Zeng N, Mariotti A, Wetzel P (2005) Terrestrial mechanisms of interannual CO<sub>2</sub> variability. *Global Biogeochem. Cycles*, **19**, GB1016, 15 pp, doi: 10.1029/2004GB002273.

## Supporting Information

Additional Supporting Information may be found in the online version of this article:

**Data S1.** Materials and Methods: National GHG Inventories

**Data S2.** Comparison with Model-based Estimates

**Data S3.** Uncertainties and Data Gaps

**Data S4.** Supporting Information Tables and Figures

**Data S5.** Supporting Information References

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.