

Decoupling and Balancing of Space and Time Errors in the Material Point Method

Mike Steffen

Martin Berzins, Mike Kirby
University of Utah

Scientific Computing and Imaging Institute

Fifth Material Point Method Workshop
Oregon State University
April 2nd, 2009

Outline

- Overview of Errors
- Observations and Motivation
- Moving Mesh MPM
- Balancing Space & Time Errors
- Sample Problem
 - Time Integration, Jump, and Quadrature Errors

Overview of Errors in MPM

- MPM simulations exhibit a variety of errors, e.g.:

Spatial	Temporal	Mixed
Approximation (h, ϕ)	Time-Integration (Δt)	Jump Error ($\Delta t, \Delta x, t$)
Mass Lumping (h, ϕ)		Quadrature ($\Delta x, t$)
		Geometric ($\Delta x, h, t$)

- MPM errors are not separable due to feedback
- Previously focused on spatial errors at one time-step
- Analysis of temporal errors is difficult due to feedback

Observations and Motivation

- Centered difference time-stepping scheme:

$$v^{k+1/2} = v^{k-1/2} + a^k \Delta t$$


$$u^{k+1} = u^k + v^{k+1/2} \Delta t$$

- “Formally second-order” in time for both velocity and displacement – assuming acceleration is well behaved
- MPM has spatial errors in acceleration:

$$v^{k+1/2} = v^{k-1/2} + \Delta t(a^k + c_1 \Delta x^2 + \dots) + c_2 \Delta t^3 + \dots$$

Moving Mesh MPM

- Moving mesh MPM
 - Mesh and material points move together
 - Can exhibit FEM mesh problems
 - Good for smooth test problems
 - Used in the MPM community
 - Particles stay in “nice” positions, quadrature jump errors no longer appear
 - Allows detailed analysis of temporal errors



Standard vs. Moving Mesh


Observations

$$v^{k+1/2} = v^{k-1/2} + \Delta t(a^k + c_1 \Delta x^2 + \dots) + c_2 \Delta t^3 + \dots$$

- Second order temporal convergence has technically not been shown in MPM
- With moving mesh MPM, and measuring local truncation errors, we can demonstrate second order temporal convergence
- By finding constants c_1 and c_2 , we can determine where spatial and temporal errors are balanced: $c_1 \Delta x^2 = c_2 \Delta t^2$


Balancing Space and Time Errors

$$v^{k+1/2} = v^{k-1/2} + \Delta t(a^k + c_1 \Delta x^2 + \dots) + c_2 \Delta t^3 + \dots$$



Spatial Acceleration Error

$$c_1 = \frac{E_a}{h^2} = 2.71 \times 10^4$$




Temporal Velocity Error

$$c_2 = \frac{E_v}{\Delta t^2} = 1.13 \times 10^7$$

Balanced when: $\Delta t = h \sqrt{\frac{c_1}{c_2}} = 2.38 \times 10^{-5}$ for 2k grid cells

Balancing Space and Time Errors

$$v^{k+1/2} = v^{k-1/2} + \Delta t(a^k + c_1 \Delta x^2 + \dots) + c_2 \Delta t^3 + \dots$$



Estimate:

$$\Delta t = 2.38 \times 10^{-5}$$

- Actual balance point (transition of convergence rate) occurs at estimate

Sample Problem

$$v_p^k = v(x_p^k) + v_{\text{ext}}^k$$



$$a = \int_0^1 f(x) dx = 0$$

$$v_{\text{ext}}^{k+1} = v_{\text{ext}}^k + a_p^k \Delta t$$

$$a_p^k = \int_{\Omega - \Omega_p^k} f(x) dx + f(x_p^k) \Delta x$$

$$x_p^{k+1} = x_p^k + v_p^k \Delta t$$

- Simpler Problem – exhibits similar errors as MPM
- Notice: $dx/dt = v(x(t))$. Analytic solution exists for piecewise-linear $v(x)$.
- External acceleration should be 0
- Quadrature errors in acceleration calculation will affect time-stepping errors



Analytic Solution

$$v_{\text{ext}}^{k+1} = v_{\text{ext}}^k + a_p^k \Delta t$$


$$v_p^0 = 0$$

$$v_p^k = v(x_p^k) + v_{\text{ext}}^k$$

$$x_p^0 = 0$$

$$x_p^{k+1} = x_p^k + v_p^k \Delta t$$

$$a_p^k = 0$$




Analytic Solution (on $0 < x < 1/3$):

$$x(t) = \frac{b + ax_0}{ae^{at_0}} e^{at} - \frac{b}{a} \quad 0 \leq t < t_{\text{cross}_1}$$

$$t_{\text{cross}_1} = \frac{1}{a} \ln \left[\frac{x_{\text{cross}} + b/a}{b + ax_0} ae^{at_0} \right]$$

Other intervals are calculated similarly



Analytic Solution

Euler Forward Error

$$v_{\text{ext}}^{k+1} = v_{\text{ext}}^k + a_p^k \Delta t$$

$$v_p^0 = 0$$

$$v_p^k = v(x_p^k) + v_{\text{ext}}^k$$

$$x_p^0 = 0$$

$$x_p^{k+1} = x_p^k + v_p^k \Delta t$$

$$a_p^k = 0$$



- Exact Spatial Integration

- Calculate Error:

$$\mathcal{E}_{EF} = x_p^k - x(t^k)$$

- Errors are $O(\Delta t)$

$$\mathcal{E}_{EF} \leq \frac{T}{2} a_{\text{max}} \Delta t$$



Centered Difference Error

$$v_{\text{ext}}^{k+1} = v_{\text{ext}}^k + a_p^k \Delta t$$

$$v_p^0 = 0$$

$$v_p^k = v(x(t^{k+1/2})) + v_{\text{ext}}^k$$

$$x_p^0 = 0$$



$$x_p^{k+1} = x_p^k + v_p^k \Delta t$$

$$a_p^k = 0$$

- Exact Spatial Integration
- Calculate Error:

$$\mathcal{E}_{EF} = x_p^k - x(t^k)$$


- Errors are $O(\Delta t^2)$



ODE Jump Error

- Different than the quadrature related grid crossing error treated in Steffen, et al.*
- Caused by integrating past spatial discontinuity in v'

One Step vs.
Two Step Method:



*M. Steffen, R. M. Kirby, M. Berzins. "Analysis and reduction of quadrature errors in the material point method (MPM)." *IJNME*, 2008


ODE Jump Error

- Tran, et al., report this as a first-order temporal error in the MPM framework:*

$$\mathcal{E}_{\text{jump}} = \bar{x}_p^{k+1} - x_p^{k+1} = (v_i^{k+1} - v_{i-1}^{k+1}) \left[\frac{x_i - x_p^k}{x_i - x_{i-1}} \Delta t_2 + \left[a_{i-1}^k + \frac{x_p^k - x_{i-1}}{x_i - x_{i-1}} (a_i^k - a_{i-1}^k) \right] \Delta t_1 \Delta t_2 \right]$$

The term: $(v_i^{k+1} - v_{i-1}^{k+1}) \left[\frac{x_i - x_p^k}{x_i - x_{i-1}} \Delta t_2 \right]$

looks to be clearly first-order in time. It *is* when x_p^k is not dependent on Δt



*L.T. Tran, J. Kim, M. Berzins, "Solving Time-Dependent PDEs using the Material Point Method, A Case Study from Gas Dynamics", IJNMF, 2009

ODE Jump Error

$$\mathcal{E}_{\text{jump}} = \bar{x}_p^{k+1} - x_p^{k+1} = (v_i^{k+1} - v_{i-1}^{k+1}) \left[\frac{x_i - x_p^k}{x_i - x_{i-1}} \Delta t_2 + \left[a_{i-1}^k + \frac{x_p^k - x_{i-1}^k}{x_i - x_{i-1}} (a_i^k - a_{i-1}^k) \right] \Delta t_1 \Delta t_2 \right]$$

Rewriting: $(v_i^{k+1} - v_{i-1}^{k+1}) \left[\frac{x_i - x_p^k}{x_i - x_{i-1}} \Delta t_2 \right]$ assuming: $x_i = x_p^k + v_p \Delta t_1$ $\Delta t_1 \leq \Delta t$

gives: $\frac{v_i^{k+1} - v_{i-1}^{k+1}}{x_i - x_{i-1}} (x_i - x_p^k) \Delta t_2 \approx \frac{\partial v}{\partial x_-} v_p \Delta t_1 \Delta t_2$

Therefore: $\mathcal{E}_{\text{jump}} = \left[\frac{\partial v}{\partial x_-} v_p + a_p \right] \Delta t_1 \Delta t_2$

The ODE jump error is then bounded by:

$$\mathcal{E}_{\text{jump}} \leq \left[\frac{\partial v}{\partial x_-} v_p + a_p \right] \frac{\Delta t^2}{4}$$

If: $\Delta t_1 = \alpha \Delta t$ $0 < \alpha < 1$
 $\Delta t_2 = (1 - \alpha) \Delta t$


Then: $\Delta t_1 \Delta t_2 = \alpha(1 - \alpha) \Delta t^2$

Therefore: $\Delta t_1 \Delta t_2 \leq \frac{1}{4} \Delta t^2$


ODE Jump Error

- Calculated: $\mathcal{E}_{jump}^k = x_p^k - \overline{x}_p^k$
- Error Bound: $\mathcal{E}_{jump} \leq \left[\frac{\partial v}{\partial x_-} v_p + a_p \right] \frac{\Delta t^2}{4}$

Jump	Jump Bound	Calculated Jump
1	2.34×10^{-8}	1.81×10^{-8}
2	9.36×10^{-8}	7.60×10^{-8}
3	2.81×10^{-7}	1.84×10^{-7}



Multiple Time Step Sizes



Single Time Step Size

Quadrature Error

- Integrating Piecewise-Constant Functions with the Midpoint Rule:*

Bounds for each time-step: $E_{\text{MAX}} = \frac{1}{2} [[f(x)]] \Delta x$

One Time-Step:

$$\begin{aligned} v^{k+1} &= v^k + (a^k + C\Delta x)\Delta t \\ x^{k+1} &= x^k + v^{k+1}\Delta t \\ &= x^k + v^k\Delta t + a^k\Delta t^2 + C\Delta x\Delta t^2 \end{aligned}$$

N Time-Steps:


$$x^N = x^0 + Nv^0\Delta t + \sum_{j=1}^N \sum_{i=1}^j a^i\Delta t^2 + \frac{1}{2}TC\Delta x\Delta t + \frac{1}{2}T^2C\Delta x$$

*M. Steffen, R.M. Kirby, M. Berzins, "Analysis and Reduction of Quadrature Errors in the Material Point Method (MPM)", *IJNME*, 2008

Quadrature Error – Final T

Effect of Time-Step

- Integration error in acceleration – effect on particle position errors:





Conclusions

- Spatial and temporal errors in MPM are interconnected, making analysis difficult.
- Moving-Mesh MPM has allows these errors to remain independent – no feedback errors
- Analysis of local truncation errors allows us to operate in unstable regimes – demonstrating expected temporal error behaviors
- Simplified problem demonstrates errors similar to MPM – easier to analyze. Hope to extend to the full MPM framework.

Acknowledgments

- Center for the Simulation of Accidental Fires and Explosions (CSAFE) DOE Grant W-7405-ENG-48



Future Work

- Extended Error analysis of simplified problem to full MPM framework.
- Through extrapolation techniques, we hope to achieve $> O(h^2)$
 - Should be easy for Moving Mesh MPM


Conclusions

- Approximation Error, Mass Lumping Error, Quadrature Error, Time integration Error, Jump Error, Feedback
- Moving Mesh MPM
 - Approximation: $O(h^2)$
 - Mass Lumping Error: $O(h^2)$
 - Quadrature Error: $O(dx^2)$
 - Time integration: $O(dt^2)$
 - No Jump Error, No Feedback.
 - Should give $O(h^2, dt^2)$ errors
- Standard MPM
 - Quadrature, Time-Stepping, Jump Error interplay
 - Interplay forces things to $O(dt)$

Quadrature Error – Final T

Effect of Particle Width

- Integration error in acceleration – effect on particle position errors:



Reminder, for $\Delta t = \Delta t_0 / 8$:

$$\mathcal{E}_{\text{ODE}} \approx 1 \cdot 10^{-3}$$

$$\mathcal{E}_{\text{jump}} \approx 4 \cdot 10^{-7}$$